A note on the minimum reduced reciprocal Randić index of n-vertex unicyclic graphs

AKBAR ALI[†,‡,*] & AKHLAQ AHMAD BHATTI[†]

[†]Department of Sciences & Humanities, National University of Computer & Emerging Sciences, B-Block, Faisal Town, Lahore-Pakistan.

[‡]Department of Mathematics, University of Gujrat, Gujrat-Pakistan.

[*]Corresponding author: akbarali.maths@gmail.com

arXiv:1601.04044v1 [math.CO] 15 Jan 2016

ABSTRACT

The graph having the minimum reduced reciprocal Randić index is characterized among the class of all unicyclic graphs with fixed number of vertices.

Keywords: Topological index; reduced reciprocal Randić index; unicyclic graph.

INTRODUCTION

All the graphs considered in the present study are simple, finite, undirected and connected. The vertex set and edge set of a graph G will be denoted by $V(G)$ and $E(G)$ respectively. The degree of a vertex $u \in V(G)$ and the edge connecting the vertices u and v will be denoted by d_u and uv respectively. Undefined notations and terminologies from (chemical) graph theory can be found in (Harary, 1969; Trinajstić, 1992).

Topological indices are numerical parameters of a graph which are invariant under graph isomorphisms. Randić (1975) proposed the following topological index:

$$R(G) = \sum_{uv \in E(G)} (d_u d_v)^{-\frac{1}{2}}$$

for measuring the extent of branching of the carbon-atom skeleton of saturated hydrocarbons and he named it as *branching index*. Nowadays, this topological index is also known as *connectivity index* and the *Randić index*. According to Gutman (2013), "the Randić index is the most investigated, most often applied, and most popular among all topological indices. Hundreds of papers and a few books are devoted to this topological index".

On the other hand, many physico-chemical properties of chemical structures are dependent on the factors different from branching. In order to take these factors into account, Estrada *et al.* (1998) introduced a modified version of the Randić index and called it as *atom-bond connectivity (ABC)* index. This index is defined as:

$$ABC(G) = \sum_{uv \in E(G)} \sqrt{\frac{d_u + d_v - 2}{d_u d_v}}.$$

Details about the chemical applicability and mathematical properties of this index can be found in the survey (Gutman, 2013), recent papers (Ahmadi *et al.*, 2014; Dimitrov, 2014; Goubko *et al.*, 2015; Palcios, 2014; Raza *et al.*, 2015) and the references cited therein.

Inspired by work on the ABC index Furtula, Graovac & Vukičević (2010) gave the following modified version of the ABC index (and hence a modified version of Randić index) under the name *augmented Zagreb index* (AZI):

$$AZI(G) = \sum_{uv \in E(G)} \left(\frac{d_u d_v}{d_u + d_v - 2} \right)^3.$$

The prediction power of AZI is better than ABC index in the study of heat of formation for heptanes and octanes (Furtula *et al.*, 2010). Details about this index can be found in the survey (Gutman, 2013), recent papers (Ali, Bhatti & Raza, 2016; Ali, Raza & Bhatti, 2016; Huang & Liu, 2015; Zhan *et al.*, 2015) and the references cited therein.

In (Manso *et al.*, 2012), a new topological index (namely Fi index) was proposed to predict the normal boiling point temperatures of hydrocarbons. In the mathematical definition of Fi index two terms are present. Gutman, Furtula & Elphick (2014), recently considered one of these terms which is given below:

$$RRR(G) = \sum_{uv \in E(G)} \sqrt{(d_u - 1)(d_v - 1)},$$

and they named it as *reduced reciprocal Randić* (RRR) *index*. In the current study, we are concerned with this recently introduced modified version of the Randić index. In order to get some preliminary information on whether this index possess any potential applicability in chemistry (especially in QSPR/QSAR studies) Gutman, Furtula & Elphick (2014) tested the correlating ability of several well known degree based topological indices (along with RRR index) for the case of standard heats (enthalpy) of formation and normal boiling points of octane isomers, and they concluded that the AZI and RRR index has the best and second-best (respectively) correlating ability among the examined topological indices (it is worth mentioning here that, among the examined topological indices ABC was also included which was the second-best degree based topological index according to the earlier study (Gutman & Tošović, 2013)). Hence it is meaningful to study the mathematical properties of the RRR index, especially bounds and characterization of the extremal elements for different graph classes. In (Gutman *et al.*, 2014), the structure of n-vertex tree having maximum RRR index and the extremal n-vertex graphs with respect to RRR index were reported.

An n-vertex (connected) graph G is unicyclic if it has n edges. Some extremal results for the unicyclic graphs can be found in the papers (Gan *et al.*, 2011; Gao & Lu, 2005; Pan *et al.*, 2006; Zhan *et al.*, 2015). The main purpose of the present note is to characterize the n-vertex unicyclic graph having minimum RRR index over the collection of all n-vertex unicyclic graphs.

MAIN RESULT

Denote by S_n^+ the unique unicyclic graph obtained from the star graph S_n by adding an edge between any two pendent vertices. Many topological indices (e.g. ABC index, R index, AZI etc.) which have S_n as an extremal graph over the set of all n-vertex trees, have also S_n^+ as an extremal graph over the set of all n-vertex unicyclic graphs. However, different

approaches required to prove these results. From the definition of RRR index, it can be easily seen that $RRR(T_n) \geq RRR(S_n) = 0$ where T_n is any n-vertex tree. Is it true that the graph S_n^+ has the minimum RRR index over the set of all n-vertex unicyclic graphs? The answer is not positive; for the n-vertex unicyclic graph H_n^+ (depicted in Fig. 1(b)), one have $RRR(H_n^+) = 1 + \sqrt{2}(2 + \sqrt{n-4})$ but on the other hand $RRR(S_n^+) = 1 + 2\sqrt{n-2}$ and

$$RRR(S_n^+) \begin{cases} < RRR(H_n^+) & \text{if } \ 5 \leq n \leq 19, \\ = RRR(H_n^+) & \text{if } \ n = 20, \\ > RRR(H_n^+) & \text{if } \ n \geq 21. \end{cases}$$

Fig. 1: (a) The n-vertex unicyclic graph H_n where n is at least 6. (b) The n-vertex unicyclic graph H_n^+ where n is at least 5.

In the following theorem we characterize the n-vertex unicyclic graph having minimum RRR index over the collection of all n-vertex unicyclic graphs for $n \geq 4$.

Theorem 1. *For any unicyclic graph U_n where $n \geq 4$, the following inequalities hold:*

$$RRR(U_n) \begin{cases} \geq 1 + 2\sqrt{n-2} & \text{if } \ 4 \leq n \leq 16, \\ \geq 1 + 3\sqrt{2} + \sqrt{n-5} & \text{if } \ n \geq 17. \end{cases}$$

The first equality holds if and only if $U_n \cong S_n^+$ and the second equality holds if and only if $U_n \cong H_n$ (where H_n is shown in Fig. 1(a)).

Proof. Routine computation yields $RRR(C_n) = n > 1 + 2\sqrt{n-2}$ for all $n \geq 4$ and $RRR(C_n) = n > 1 + 3\sqrt{2} + \sqrt{n-5}$ for all $n \geq 7$, so we assume $U_n \ncong C_n$. Let $P(U_n) = \{u_0, u_1, u_2, ..., u_{p-1}\}$ be the set of all pendent vertices in U_n. For $1 \leq i \leq p-1$, suppose that v_i is adjacent with u_i and W_{u_i} is the set of all pendent neighbors of v_i different from u_i. Choose a member of $P(U_n)$, say u_0 (without loss of generality), such that

1. the number of elements in W_{u_0} is as large as possible;

2. subject to (1), d_{v_0} is as small as possible.

Let $d_{v_0} = x$ and $N(v_0) = \{u_0, u_1, u_2, ..., u_{p-1}, u_p, ..., u_{x-1}\}$ where $d_{u_i} = 1$ for $0 \leq i \leq p-1$ and $d_{u_i} \geq 2$ for $p \leq i \leq x-1$ (see Fig. 2).



Fig. 2: The presentation of an n-vertex unicyclic graph U_n used in the proof of Theorem 1.

If U'_{n-1} is the graph obtained from U_n by removing the vertex u_0, then

$$RRR(U_n) = RRR(U'_{n-1}) + \sum_{i=1}^{x-1} \left[\sqrt{(x-1)(d_{u_i}-1)} - \sqrt{(x-2)(d_{u_i}-1)} \right] \tag{1}$$

We will discuss three cases:

Case 1. Either the vertex v_0 is adjacent with at least two non-pendent vertices or v_0 is adjacent with exactly one non-pendent vertex u_{x-1} such that $d_{u_{x-1}} \geq 5$ (that is either $p \leq x-2$ or $p = x-1, d_{u_{x-1}} \geq 5$).

Let $\mathcal{U}_n^{(1)}$ be the collection of all those n-vertex unicyclic graphs (different from C_n) which fall in this case. By using induction on n, we will prove that the only one graph, namely S_n^+, has the minimum RRR value among all the members of $\mathcal{U}_n^{(1)}$. [Then the desired result will follow from the fact that

$$RRR(S_n^+) = 1 + 2\sqrt{n-2} > 1 + 3\sqrt{2} + \sqrt{n-5} = RRR(H_n) \quad \text{for all } n \geq 17.]$$

For $n = 4$, there are only two non-isomorphic unicyclic graphs namely C_n and S_n^+ and hence the result holds for $n = 4$. For $n = 5$, all the non-isomorphic members of $\mathcal{U}_n^{(1)}$ are depicted in the Fig. 3 along with their RRR values.



Fig. 3: All the non-isomorphic members of $\mathcal{U}_5^{(1)}$ together with their RRR values.

Now, suppose that $U_n \in \mathcal{U}_n^{(1)}$ and $n \geq 6$. By virtue of inductive hypothesis and from Equation (1), one have

$$RRR(U_n) \geq 1 + 2\sqrt{n-3} + \left(\sqrt{x-1} - \sqrt{x-2}\right) \sum_{i=1}^{x-1} \sqrt{d_{u_i}-1}, \tag{2}$$

with equality if and only if $U'_{n-1} \cong S^+_{n-1}$. We discuss two subcases:

Subcase 1.1. If $p \geq 2$. Then from Inequality (2) it follows that

$$RRR(U_n) \geq 1 + 2\sqrt{n-3} + \left(\sqrt{x-1} - \sqrt{x-2}\right) \sum_{i=p}^{x-1} \sqrt{d_{u_i} - 1} \tag{3}$$

According to the definition of $U_n \in \mathcal{U}_n^{(1)}$, either $p \leq x-2$ or $p = x-1, d_{u_{x-1}} \geq 5$. If $p \leq x-2$, then Inequality (3) implies that

$$\begin{aligned} RRR(U_n) &\geq 1 + 2\sqrt{n-3} + \left(\sqrt{x-1} - \sqrt{x-2}\,\right)(x-p) \\ &\geq 1 + 2\sqrt{n-3} + 2\left(\sqrt{x-1} - \sqrt{x-2}\,\right) \\ &\geq 1 + 2\sqrt{n-3} + 2\left(\sqrt{n-2} - \sqrt{n-3}\,\right) = 1 + 2\sqrt{n-2}. \end{aligned}$$

The equality $RRR(U_n) = 1 + 2\sqrt{n-2}$ holds if and only if $x = n-1, x-p = 2$ and $U'_{n-1} \cong S^+_{n-1}$.

If $p = x-1$ and $d_{u_{x-1}} \geq 5$, then the graph U'_{n-1} must be different from S^+_{n-1} and hence from Equation (1), one have

$$\begin{aligned} RRR(U_n) &> 1 + 2\sqrt{n-3} + 2\left(\sqrt{x-1} - \sqrt{x-2}\,\right) \\ &> 1 + 2\sqrt{n-3} + 2\left(\sqrt{n-3} - \sqrt{n-4}\,\right) \quad \text{(since in this case } x < n-2\text{)} \\ &> 1 + 2\sqrt{n-2}. \end{aligned}$$

Subcase 1.2. If $p = 1$. Then, from the definition of u_0, it follows that the set W_{u_i} is empty for all $u_i \in P(U_n)$. It means that no two pendent edges are adjacent.

If $x \geq 4$, then among the vertices $u_1, u_2, ..., u_{x-1}$ at least two are disconnected in $U_n - v_0$ (because otherwise U_n contains more than one cycle, a contradiction. See Fig. 4 for the graphs $U_n - v_0$ and U_n considered in this subcase).



Fig. 4: The graphs U_n and $U_n - v_0$ used in Subcase 1.2 of the proof of Theorem 1.

Without loss of generality, let u_1 and u_2 are disconnected in $U_n - v_0$ and suppose that C_i (for $i = 1, 2$) is the component of $U_n - v_0$ containing u_i (for $i = 1, 2$). Since $d_{u_i} \geq 2$ in U_n for all i $(1 \leq i \leq x-1)$, so both the components C_1 and C_2 must be non-trivial. Note that

at least one of C_1 and C_2 must be a tree (for otherwise U_n contains more than one cycle, a contradiction). Let C_1 is a tree. Since every non-trivial tree contains at least two pendent vertices and no two pendent edges of U_n are adjacent, so there exist $w_1 \in V(C_1) \cap P(U_n)$ and $w_1w_2 \in V(C_1) \cap E(U_n)$ such that $d_{w_2} = 2$ in U_n, which contradicts the definition of u_0. Hence $x = 2$ or 3. It should be noted that the graph U'_{n-1} is different from S^+_{n-1} in this case. Now, we consider further two subcases:

Subcase 1.2.1. If $x = 2$. Then from the Inequality (1), we have

$$\begin{aligned} RRR(U_n) &> 1 + 2\sqrt{n-3} + \sqrt{d_{u_1} - 1} \\ &\geq 3 + 2\sqrt{n-3} > 1 + 2\sqrt{n-2}. \end{aligned}$$

Subcase 1.2.2. If $x = 3$. Then from the Inequality (1), it follows that

$$\begin{aligned} RRR(U_n) &> 1 + 2\sqrt{n-3} + \left(\sqrt{2} - 1\right)\left(\sqrt{d_{u_1} - 1} + \sqrt{d_{u_2} - 1}\ \right) \\ &\geq 1 + 2\sqrt{n-3} + 2\left(\sqrt{2} - 1\right) \\ &> 1 + 2\sqrt{n-2}, \quad \text{because } n \geq 6. \end{aligned}$$

Therefore, for any $U_n \in \mathcal{U}^{(1)}_n$ we have $RRR(U_n) \geq RRR(S^+_n)$ with equality if and only if $U_n \cong S^+_n$.

Case 2. The vertex v_0 is adjacent with exactly one non-pendent vertex u_{x-1} such that $d_{u_{x-1}} = 3$ or 4 (that is $p = x - 1$ and $d_{u_{x-1}} = 3$ or 4).

Let $\mathcal{U}^{(2)}_n$ be the family of all those n-vertex unicyclic graphs (different from C_n) which fall in this case. Note that n must be at least five in this case. By using induction on n, we will prove that the only one graph, namely H^+_n, has the minimum RRR value among all the members of $\mathcal{U}^{(2)}_n$. [Then the desired result will follow from the fact that

$$RRR(H^+_n) = 1 + 2\sqrt{2} + \sqrt{2(n-4)} > \begin{cases} 1 + 2\sqrt{n-2} & \text{for } 5 \leq n \leq 16, \\ 1 + 3\sqrt{2} + \sqrt{n-5} & \text{for } n \geq 17.] \end{cases}$$

It can be easily seen that $\mathcal{U}^{(2)}_5$ has only one element namely H^+_5. For $n = 6$, all the non-isomorphic members of $\mathcal{U}^{(2)}_n$ are depicted in the Fig. 5 along with their RRR values.



Fig. 5: All the non-isomorphic members of $\mathcal{U}^{(2)}_6$ together with their RRR values.

Hence the result holds for $n = 5, 6$. Suppose that $U_n \in \mathcal{U}_n^{(2)}$ and $n \geq 7$. By using the inductive hypothesis in the Equation (1), we have

$$RRR(U_n) \geq 1 + 2\sqrt{2} + \sqrt{2(n-5)} + \left(\sqrt{x-1} - \sqrt{x-2}\right) \sum_{i=1}^{x-1} \sqrt{d_{u_i} - 1}, \tag{4}$$

with equality if and only if $U'_{n-1} \cong H^+_{n-1}$. We consider two subcases:

Subcase 2.1. If $p \geq 2$. Then from Inequality (4) it follows that

$$RRR(U_n) \geq 1 + 2\sqrt{2} + \sqrt{2(n-5)} + \left(\sqrt{x-1} - \sqrt{x-2}\right) \sum_{i=p}^{x-1} \sqrt{d_{u_i} - 1} \tag{5}$$

According to the definition of $U_n \in \mathcal{U}_n^{(2)}$, $p = x - 1$ and $d_{u_{x-1}} = 3$ or 4. Hence from Equation (5), it follows that

$$\begin{aligned} RRR(U_n) &\geq 1 + 2\sqrt{2} + \sqrt{2(n-5)} + \left(\sqrt{x-1} - \sqrt{x-2}\,\right) \sqrt{d_{u_{x-1}} - 1} \\ &\geq 1 + 2\sqrt{2} + \sqrt{2(n-5)} + \sqrt{2}\left(\sqrt{x-1} - \sqrt{x-2}\,\right) \\ &\geq 1 + 2\sqrt{2} + \sqrt{2(n-5)} + \sqrt{2}\left(\sqrt{n-4} - \sqrt{n-5}\,\right) \text{ (since } x \leq n-3) \\ &= 1 + 2\sqrt{2} + \sqrt{2(n-4)}. \end{aligned}$$

The equality $RRR(U_n) = 1 + 2\sqrt{2} + \sqrt{2(n-4)}$ holds if and only if $x = n - 3, d_{u_{x-1}} = 3$ and $U'_{n-1} \cong H^+_{n-1}$.

Subcase 2.2. If $p = 1$. Then, $x = 2$ because $p = x - 1$. It is easy to see that the graph U'_{n-1} is different from H^+_{n-1} in this case. From the Inequality (1), we have

$$\begin{aligned} RRR(U_n) &> 1 + 2\sqrt{2} + \sqrt{2(n-5)} + \sqrt{d_{u_1} - 1} \\ &\geq 1 + 3\sqrt{2} + \sqrt{2(n-5)} > 1 + 2\sqrt{2} + \sqrt{2(n-4)}. \end{aligned}$$

Therefore, for any $U_n \in \mathcal{U}_n^{(2)}$ we have $RRR(U_n) \geq RRR(H^+_n)$ with equality if and only if $U_n \cong H^+_n$.

Case 3. The vertex v_0 is adjacent with exactly one non-pendent vertex u_{x-1} such that $d_{u_{x-1}} = 2$ (that is $p = x - 1$ and $d_{u_{x-1}} = 2$).

Let $\mathcal{U}_n^{(3)}$ be the class of all those n-vertex unicyclic graphs (different from C_n) which fall in this case. Note that n must be at least 6 in this case. By using induction on n, we will prove that the only one graph, namely H_n, has the minimum RRR value among all the members of $\mathcal{U}_n^{(3)}$. [Then the desired result will follow from the fact that

$$RRR(H_n) = 1 + 3\sqrt{2} + \sqrt{n-5} > 1 + 2\sqrt{n-2} \quad \text{for} \quad 6 \leq n \leq 16.]$$

It can be easily seen that $\mathcal{U}_6^{(3)}$ has only one member, namely H_6. For $n = 7$, all the non-isomorphic members of $\mathcal{U}_n^{(3)}$ are depicted in the Fig.6 along with their RRR values.



Fig. 6: All the non-isomorphic members of $\mathcal{U}_7^{(3)}$ together with their RRR values.

Hence the result holds for $n = 6, 7$. Suppose that $U_n \in \mathcal{U}_n^{(3)}$ and $n \geq 8$. By virtue of inductive hypothesis and from Equation (1), one have

$$RRR(U_n) \geq 1 + 3\sqrt{2} + \sqrt{n-6} + \left(\sqrt{x-1} - \sqrt{x-2}\right) \sum_{i=1}^{x-1} \sqrt{d_{u_i} - 1}, \tag{6}$$

with equality if and only if $U'_{n-1} \cong H_{n-1}$. We consider two subcases:

Subcase 3.1. If $p \geq 2$. Then from Inequality (6) it follows that

$$RRR(U_n) \geq 1 + 3\sqrt{2} + \sqrt{n-6} + \left(\sqrt{x-1} - \sqrt{x-2}\right) \sum_{i=p}^{x-1} \sqrt{d_{u_i} - 1} \tag{7}$$

By the definition of $U_n \in \mathcal{U}_n^{(3)}$, $p = x - 1$ and $d_{u_{x-1}} = 2$. Hence from Equation (7), it follows that

$$\begin{aligned} RRR(U_n) &\geq 1 + 3\sqrt{2} + \sqrt{n-6} + \sqrt{x-1} - \sqrt{x-2} \\ &\geq 1 + 3\sqrt{2} + \sqrt{n-5}. \end{aligned}$$

The last inequality holds because $x \leq n - 4$. Furthermore, the equality $RRR(U_n) = 1 + 3\sqrt{2} + \sqrt{n-5}$ holds if and only if $x = n - 4$ and $U'_{n-1} \cong H_{n-1}$.

Subcase 3.2. If $p = 1$. Then, $x = 2$ because $p = x - 1$. Observe that the graph U'_{n-1} is different from H_{n-1} in this case. From the Inequality (1), we have

$$\begin{aligned} RRR(U_n) &> 1 + 3\sqrt{2} + \sqrt{n-6} + \sqrt{d_{u_1} - 1} \\ &= 2 + 3\sqrt{2} + \sqrt{n-6} > 1 + 3\sqrt{2} + \sqrt{n-5}. \end{aligned}$$

Therefore, for any $U_n \in \mathcal{U}_n^{(3)}$ we conclude that $RRR(U_n) \geq RRR(H_n)$ with equality if and only if $U_n \cong H_n$. This completes the proof.

REFERENCES

Ahmadi, M. B., Dimitrov, D., Gutman, I. & Hosseini, S. A. 2014. Disproving a conjecture on trees with minimal atom-bond connectivity index. MATCH Communications in Mathematical and in Computer Chemistry **72**: 685-698.

Ali, A., Bhatti, A. A. & Raza, Z. 2016 (in press). The augmented Zagreb index, vertex connectivity and matching number of graphs. Bulletin of the Iranian Mathematical Society.

Ali, A., Raza, Z. & Bhatti, A. A. 2016 (in press). On the augmented Zagreb index. Kuwait Journal of Science.

Dimitrov, D. 2014. On structural properties of trees with minimal atom-bond connectivity index. Discrete Applied Mathematics **172**: 28-44.

Estrada, E., Torres, L., Rodrguez, L. & Gutman, I. 1998. An atom-bond connectivity index: modelling the enthalpy of formation of alkanes. Indian Journal of Chemistry-Section A **37**: 849-855.

Furtula, B., Graovac, A. & Vukičević, D. 2010. Augmented Zagreb index. Journal of Mathematical Chemistry **48**: 370-380.

Gan, L., Hou, H. & Liu, B. 2011. Some results on atom-bond connectivity index of graphs. MATCH Communications in Mathematical and in Computer Chemistry **66**: 669-680.

Gao, J. & Lu, M. 2005. On the Randić index of unicyclic graphs. MATCH Communications in Mathematical and in Computer Chemistry **53**: 377-384.

Goubko, M., Magnant, C., Nowbandegani, P. S. & Gutman, I. 2015. ABC index of trees with fixed number of leaves. MATCH Communications in Mathematical and in Computer Chemistry **74**: 697-702.

Gutman, I. 2013. Degree-based topological indices. Croatica Chemica Acta **86**: 351-361.

Furtula, B., Gutman, I. & Elphick, C. 2014. Three new/old vertex-degree-based topological indices. MATCH Communications in Mathematical and in Computer Chemistry **72**: 617-632.

Gutman, I. & Tošović, J. 2013. Testing the quality of molecular structure descriptors: Vertex-degree-based topological indices. Journal of the Serbian Chemical Society **78**: 805-810.

Harary, F. 1969. Graph Theory. Addison-Wesley, Reading, MA.

Huang, Y. & Liu, B. 2015. Ordering graphs by the augmented Zagreb indices. Journal of Mathematical Research with Applications **35**(2): 119-129.

Manso, F. C. G., Júnior, H. S., Bruns, R. E., Rubira, A. F. & Muniz, E. C. 2012. Development of a new topological index for the prediction of normal boiling point temperatures of hydrocarbons: The Fi index. Journal of Molecular Liquids **165**: 125-132.

Palacios, J. L. 2014. A resistive upper bound for the ABC index. MATCH Communications in Mathematical and in Computer Chemistry **72**: 709-713.

Pan, X. F., Xu, J. M. & Yang, C. 2006. On the Randić index of unicyclic graphs

with k pendant vertices. MATCH Communications in Mathematical and in Computer Chemistry **55**: 409-417.

Randić, M. 1975. On characterization of molecular branching. Journal of the American Chemical Society **97**: 6609-6615.

Raza, Z., Bhatti, A. A. & Ali, A. 2015 (in press). More on comparison between first geometric-arithmetic index and atom-bond connectivity index. Miskolc Mathematical Notes.

Trinajstić, N. 1992. Chemical Graph Theory (2nd revised ed.). CRC Press, Florida.

Zhan, F., Qiao, Y. & Cai, J. 2015. Unicyclic and bicyclic graphs with minimal augmented Zagreb index. Journal of Inequalities and Applications. DOI 10.1186/s13660-015-0651-2.